

Clinical Policy: Nerve Blocks and Neurolysis for Pain Management

Reference Number: CP.MP.170

Date of Last Revision: 12/21

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Nerve blocks are the temporary interruption of conduction of impulses in peripheral nerves or nerve trunks created by the injection of local anesthetic solutions. They can be used to identify the source of pain or to treat pain.

Note: For sacroiliac nerve block and radiofrequency neurotomy, please refer to CP.MP.166 Sacroiliac Joint Interventions

Policy/Criteria

It is the policy of health plans affiliated with Centene Corporation[®] that invasive pain management procedures performed by a physician are **medically necessary** when *the relevant criteria are met and the patient receives only one procedure per visit, with or without radiographic guidance.*

Table of Contents

I. Occipital Nerve Block.....	1
II. Sympathetic Nerve Blocks.....	2
III. Celiac Plexus Nerve Block/Neurolysis.....	3
IV. Intercostal Nerve Block/Neurolysis	3
V. Genicular Nerve Blocks and Genicular Nerve Radiofrequency Neurotomy	3
VI. Peripheral nerve blocks.....	4
VII. Intraosseous radiofrequency nerve ablation of basivertebral nerve.....	4

I. Occipital Nerve Block

A. *An initial injection* of a local anesthetic for the diagnosis of suspected occipital neuralgia is **medically necessary** when all of the following are met:

1. Patient has unilateral or bilateral pain located in the distribution of the greater, lesser and/or third occipital nerves;
2. Pain has two of the following three characteristics:
 - a. Recurring in paroxysmal attacks lasting from a few seconds to minutes;
 - b. Severe intensity;
 - c. Shooting, stabbing, or sharp in quality;
3. Pain is associated with dysaesthesia and/or allodynia apparent during innocuous stimulation of the scalp and/or hair, and at least one of the following:
 - a. Tenderness over the affected nerve branches;

- b. Trigger point at the emergence of the greater occipital nerve or in the distribution of C2.
- B.** *Therapeutic occipital nerve blocks* are **medically necessary** when all of the following are met:
1. There was temporary relief from the initial/previous injection;
 2. The member/enrollee has failed 3 months of conservative treatment including all of the following:
 - a. Heat, rest and/or physical therapy, including massage;
 - b. NSAIDS, unless contraindicated or not tolerated;
 - c. Oral anticonvulsant medications (e.g., carbamazepine, gabapentin, pregabalin) or tricyclic antidepressants;
 - d. Activity modification to address triggers;
 3. No more than 4 injections are to be given within 12 months (includes diagnostic injection).
- C.** *Occipital nerve block* for the diagnosis or treatment of other types of headaches, including migraine and cervicogenic headaches, is considered **not medically necessary**.

Note: Please refer to CP.PHAR.232 OnabotulinumtoxinA (Botox) for requests for Botox injections for migraines

II. Sympathetic Nerve Blocks have limited evidence to prove effectiveness of treatment and consideration will be made on a case by case basis. The criteria below provides a basis for documenting patient-specific clinical information to help guide clinical decision making.

- i. *First or second sympathetic nerve block*:
 1. Diagnosis of *complex regional pain syndrome* (CRPS) (also called reflex sympathetic dystrophy) and all of the following:
 - a. Pain is being managed by a pain management specialist with experience treating CRPS;
 - b. The member/enrollee is in an active rehabilitation regimen;
 - c. Failed ≥ 3 weeks of conservative therapies such as activity modification, exercises, topical capsaicin cream, and oral medical management such as nonsteroidal anti-inflammatories, antidepressants, anticonvulsants and glucocorticoids;
 - d. Two or more of the following findings of the involved digit/extremity:
 - i. Hyperalgesia or allodynia (pain sensation in response to a typically non-painful stimulus);
 - ii. Evidence of edema and/or sweating changes and/or sweating asymmetry;
 - iii. Evidence of temperature asymmetry ($>1^{\circ}\text{C}$) and/or skin color changes and/or asymmetry;
 - iv. Evidence of decreased range of motion and/or motor dysfunction (weakness, tremor, dystonia) and/or trophic changes (hair, nail, skin).

CLINICAL POLICY

Nerve Blocks

- ii. *Additional sympathetic nerve blocks for CRPS* may be considered **medically necessary** when all of the following are met:
 1. Nerve blocks are given at least a week apart;
 2. There was an immediate positive response to the first or second nerve block (eg, improved temperature and decreased pain).
- iii. *Additional sympathetic nerve blocks* without documented benefit from the first or second are **not medically necessary**.
- iv. Sympathetic nerve blocks for any other indication, including ischemic limb pain, are **not medically necessary** as there is a lack of evidence to support effectiveness.

III. Celiac Plexus Nerve Block/Neurolysis

- A. *Celiac plexus nerve block/neurolysis* is **medically necessary** for either of the following indications:
 1. Chronic neuralgic pain secondary to pancreatic cancer, all of the following:
 - a. Diagnosis of pancreatic cancer with severe visceral abdominal/back pain;
 - b. Strong analgesics such as opioids are no longer effective or their side effects decrease quality of life;
 - c. No malignancy in an area of somatic innervation such as the peritoneum or diaphragm.
 2. Refractory pain due to chronic pancreatitis with non-dilated pancreatic duct.
- B. A repeat *celiac plexus nerve block* for refractory pain from chronic pancreatitis with non-dilated pancreatic duct is **medically necessary** when both of the following are met:
 1. At least three months have passed since previous injection;
 2. There was a clinical benefit from the initial celiac block.
- C. Repeat *celiac plexus nerve blocks or neurolysis*, for any indication other than those noted above, are **not medically necessary** as there is a lack of evidence to support effectiveness.

IV. Intercostal Nerve Block/Neurolysis

- A. *Intercostal nerve block/neurolysis* is **medically necessary** for chronic neuralgic pain secondary to an injured intercostal nerve as a result of a rib fracture, a thoracotomy incision or chronic pain due to post herpetic neuralgia, or other neuropathic process when all of the following are met:
 1. Suspected organic problem;
 2. Non-responsiveness to conservative modalities of treatment;
 3. Pain and disability of moderate to severe degree;
 4. No evidence of contraindications such as infection or pain of predominately psychogenic origin.

V. Genicular Nerve Blocks, Neurolysis and Genicular Nerve Radiofrequency Neurotomy

There is insufficient evidence to determine safety and effectiveness of *genicular nerve blocks, neurolysis and radiofrequency neurotomy of the articular nerve*.

VI. Peripheral/Ganglion Nerve Blocks

Note: If administered as part of a surgery or other procedure, coding for peripheral/ganglion nerve blocks should follow proper coding practices and would not be subject to prior authorization or payment separately from the procedure.

- A. *Peripheral nerve blocks for diagnosis and treatment of malignant pain* are considered **medically necessary** as part of a comprehensive pain management program.
- B. *Peripheral nerve blocks for diagnosis or treatment of post-herniorrhaphy pain* are considered **medically necessary** when all of the following criteria are met:
 - 1. A first diagnostic peripheral nerve block when all of the following are met:
 - a. Diagnosis of post-herniorrhaphy neuralgia;
 - b. Groin pain has persisted for three months after surgical hernia repair;
 - c. Less invasive pain management methods such as NSAIDs and/or opiates have not relieved the pain;
 - d. Imaging studies have ruled out non-neuropathic causes of pain;
 - e. Documentation indicates that pain is not attributable to any other cause;
 - 2. Therapeutic peripheral nerve block(s) for treatment of post-herniorrhaphy pain when all of the following are met:
 - a. There was temporary relief from the initial/previous injection;
 - b. Injections are given at least a week apart.
- C. *Peripheral nerve blocks for prevention or treatment of headaches*, including, but not limited to: migraine headaches, treatment-refractory migraines in pregnancy, and short-lasting unilateral neuralgiform headaches, are considered **not medically necessary** as effectiveness has not been established.
- D. There is insufficient evidence in the published peer-reviewed literature to support the use of *peripheral nerve blocks for the treatment of trigeminal neuralgia*.
- E. There is insufficient evidence in the published peer-reviewed literature to support the use of *peripheral/ganglion nerve blocks or neurolysis* for any condition not indicated elsewhere in this policy, including chronic pain. There is ongoing research but insufficient evidence to establish efficacy.

VII. Intraosseous Radiofrequency Nerve Ablation of the Basivertebral Nerve

There is insufficient evidence to determine the safety and effectiveness of *intraosseous radiofrequency nerve ablation of the basivertebral nerve* (e.g. Intracept[®] Intraosseous Nerve Ablation System.) for the treatment of chronic low back pain.

Background

Local Injections for Cervicogenic and Occipital Neuralgia

Greater occipital nerve blocks have been advocated as a diagnostic test for cervicogenic headache and occipital neuralgia. The effectiveness of greater occipital nerve block in patients

Nerve Blocks

with primary headache syndromes is controversial. The International Headache Society (IHS) defines occipital neuralgia as unilateral or bilateral paroxysmal, shooting or stabbing pain in the posterior part of the scalp, in the distribution of the greater, lesser or third occipital nerves, sometimes accompanied by diminished sensation or dysaesthesia in the affected area and commonly associated with tenderness over the involved nerve(s).¹ The IHS includes relief of pain following a local anesthetic block of the affected nerve as part of their diagnostic criteria for occipital neuralgia. Thus, the principal indication for occipital block is diagnosis. Another indication is the treatment of chronic occipital neuralgia, often with a series of therapeutic blocks combining local anesthetic and corticosteroid. Pain relief is typically prompt and may last several weeks or even months. At that time the injection may be repeated.

Sympathetic Nerve Blocks

Sympathetic nerves may be injected for several reasons:

- Diagnostic - to determine the source of pain, e.g., to identify or pinpoint a nerve that acts as a pathway for pain; to determine the type of nerve that conducts the pain; to distinguish between pain that is central (within the spinal cord) or peripheral (outside the spinal cord) in origin; or to determine whether a neurolytic block or surgical lysis of the nerve should be performed;
- Therapeutic - to treat painful conditions that respond to nerve blocks (e.g., celiac block for pain of pancreatic cancer); and
- Prognostic - to predict the outcome of long-lasting interventions (e.g., lumbar sympathectomy).

The response to sympathetic blockade is the best diagnostic test for CRPS. If the patient has had a technically successful sympathetic block and does not obtain significant relief, then the patient probably does not have CRPS. Over two thirds of patients will obtain significant relief with minimal effect on motor and sensory function because the sympathetic fibers are the least myelinated (as compared to motor and sensory nerve fibers) these fibers are the first to be affected by the local anesthetic.

A 2014 case report and literature review identified only five cases, and no Level I or II evidence-based trials to support the use of sympathetic nerve block for ischemic pain.¹⁶ The authors presented two cases of patients who experienced severe pain due to ischemia despite full regional nerve blocks.¹⁶ The available literature is not sufficient to support the use of sympathetic nerve blocks for ischemic limb pain.

Celiac Plexus Nerve Block/Neurolysis for Pancreatic Cancer

Although its analgesic effectiveness is similar to analgesic drugs, celiac plexus neurolysis offers pain reduction without the significant adverse effects of opiates.² A multidisciplinary, international guideline issued a strong recommendation based on moderate quality evidence for celiac plexus neurolysis as a treatment for pain associated with advanced pancreatic cancer.² Furthermore, a 2011 Cochrane review stated that celiac plexus block (neurolysis) significantly reduced opiate use and lowered pain compared to the control group.³

The optimal timing of celiac plexus neurolysis for pain due to pancreatic cancer is not known.² Advocates of an earlier approach argue that pain is more effectively addressed by neurolysis

CLINICAL POLICY

Nerve Blocks

when treated earlier, and opiate-related side effects may also be reduced compared to later treatment. However, the effects of celiac plexus neurolysis diminish over time, which would leave a patient with fewer options as the cancer progresses and pain becomes more severe. Repeat celiac plexus neurolysis for pain due to pancreatic cancer is effective only about 30% of the time and is not recommended.^{2, 17}

Celiac Plexus Nerve Block/Neurolysis for Chronic Pancreatitis

Celiac plexus blockade is an option for pain relief in patients with refractory pain due to chronic pancreatitis and a non-dilated pancreatic duct. Advantages of celiac plexus blockade include that a single treatment can potentially provide pain reduction or relief, may reduce or eliminate the need for oral analgesia, and can be performed quickly and repeated as needed. However, it is unclear which patients will derive the most benefit and the pain relief is transient, lasting for three to six months.²⁴

The American College of Gastroenterology suggests considering celiac plexus block for treatment of pain in chronic pancreatitis (conditional recommendation, very low quality of evidence) noting that celiac plexus blockade represents a relatively low-risk, opioid-free method to reduce refractory pain in certain patients with chronic pancreatitis.⁴¹

Intercostal Nerve Blocks

Intermittent intercostal nerve blocks can be used to control pain in the chest and upper abdomen, such as pain associated with rib fractures or chronic pain due to post herpetic neuralgia. Intercostal nerve blocks can be performed using anatomic landmarks or with ultrasound guidance, which can be used to minimize the chance of intravascular injection and pneumothorax and to increase reliable dermatomal coverage.^{4, 8}

For isolated injuries, such as single rib fracture, nonsteroidal anti-inflammatory drugs with or without opioids would be the initial treatment. For more severe injuries, particularly if ventilation is compromised, intercostal nerve blocks may be needed. For patients with multiple rib fractures, there is a need to perform the procedure at multiple intercostal levels. Repeated blockade may be needed for prolonged relief upon return of pain and/or deterioration in functional status. For repeat blocks or other interventions, patient must have been responsive to prior interventions with improvement in physical and functional status.^{5, 8}

Regional anesthesia plays an important role in thoracic surgery, particularly with regard to post-operative pain control. The first choice of regional anesthesia for thoracic surgery is epidural analgesia or thoracic paravertebral block. In general, the analgesic efficiencies of both these types of anesthesia are equivalent; however, thoracic paravertebral block has some advantages over epidural analgesia, including fewer complications. When these two blocks are contraindicated, intercostal nerve block or interpleural block should be considered.^{6, 7}

Genicular Nerve Blocks and Radiofrequency Neurotomy

The genicular nerve is a sensory nerve that surrounds the knee and provides innervation for the joint. Genicular nerve blocks, neurolysis and radiofrequency neurotomy are emerging interventions for knee pain. The limited evidence regarding genicular nerve blocks for determining appropriateness of treatment with genicular radiofrequency ablation has reached

CLINICAL POLICY

Nerve Blocks

conflicting results.^{9, 10, 41} A few small studies suggest that genicular radiofrequency neurotomy may be effective for relief of pain, but further research is needed to establish safety and efficacy.
11-15

Peripheral/Ganglion Nerve Block.

Peripheral nerve blocks (PNB) are widely-used for surgical anesthesia as well as for both postoperative and nonsurgical analgesia. Indications for PNBs are diverse and vary widely. Blocks are often used to avoid the effects of alternative anesthetics or analgesics. The most common rationale for their use is to avoid side effects and complications of general anesthesia, particularly respiratory-related effects, and to provide analgesia while minimizing opioid use.³⁷

Chronic pain can be treated with a number of pharmacologic and nonpharmacologic therapies which generally fall into six major categories: pharmacologic, physical medicine, behavioral medicine neuromodulation, interventional and surgical approaches. Optimal outcomes result from multiple approaches. Interventional approaches, which typically attempt to target the presumed pain generators, may play a complementary role to other strategies (e.g., rehabilitation and appropriate pharmacotherapy.) The best candidates for interventional management have persistent focal pain of shorter duration, appropriate expectations, and well-managed psychosocial distress.³³

Cancer pain can be caused by complex interactions among cancer cells, the peripheral and central nervous systems, and the immune system. Peripheral pain receptors may become activated, sensitized or injured with certain cancers. Neuropathic pain may arise from nerve tissue damage and cancer patients may experience mild to severe pain. At least 15% will experience no relief or have severe adverse effects from interventions to address their pain. Nerve blocks or other interventional procedures may be appropriate as part of a comprehensive pain management program.^{34, 35}

Peripheral Nerve Blocks for Prevention or Treatment of Headaches

Peripheral nerve blocks have been proposed as a treatment for migraines in pregnancy and refractory migraines. However, evidence is limited to support this indication. In a series of 13 pregnant women with migraine refractory to medication, injection of local anesthetic into one or more peripheral nerve resulted in elimination of pain in seven women, pain reduction in two and no response in four women. Larger studies are necessary to better define the efficacy of this approach.³¹

Peripheral Nerve Blocks for Diagnosis and Treatment of Post-Herniorrhaphy Groin Pain

Persistent pain following inguinal hernia surgery is relatively common and a comprehensive pain management program is recommended. A prospective study, including elective primary open hernia repairs, found persistent pain occurred in 16.5-16.1 percent of patients at six months and five years.³⁶ Acute pain persisting more than eight weeks is most likely neuropathic due to primary or secondary nerve injuries. Post-herniorrhaphy neuralgia should be suspected if pain persists beyond six to eight weeks. These patients should undergo imaging to exclude nonneuropathic causes. Patients with positive response to initial nerve block can be treated every 1-3 weeks until pain relief is sustained. Those who do not obtain pain relief may require groin nerve sacrifice.³⁶

Peripheral Nerve Blocks for Prevention or Treatment of Trigeminal Neuralgia

Compression of the trigeminal nerve root is the main mechanism of trigeminal neuralgia, but brainstem lesions account for a small proportion of cases. Initial treatment of most patients with trigeminal neuralgia is pharmacologic therapy. For patients with TN refractory to medical therapy, it is reasonable to discuss options for surgical therapy (e.g., microvascular decompression, various types of rhizotomy, or gamma knife radiosurgery.) The decision to have surgery and the choice among surgical options will be influenced by individual circumstances including patient preference, adverse effect profile of the available techniques, and expertise of the local center.⁴¹ There is insufficient evidence in the published peer-reviewed literature to support the use of peripheral nerve blocks for the treatment of trigeminal neuralgia.

Intraosseous Radiofrequency Nerve Ablation of Basivertebral Nerve

Basivertebral nerve radiofrequency ablation has been developed for the treatment of chronic low back pain thought to arise from the vertebral body endplates. The Intracept Intraosseous Nerve Ablation System, Relieva Medsystems, Inc. is approved by the FDA and intended to be used in conjunction with radiofrequency generators for the ablation of basivertebral nerves of the L3 through S1 vertebrae. Its purpose is to relieve chronic low back pain of at least six months duration that has not responded to at least six months of conservative care, and is also accompanied by features consistent with Type 1 or Type 2 Modic changes on an MRI [e.g., inflammation, edema, vertebral endplate changes, disruption and fissuring of the endplate, vascularized fibrous tissues within the adjacent marrow, hypointensive signals (Type 1 Modic change), and changes to the vertebral body marrow including replacement of normal bone marrow by fat, and hyperintensive signals (Type 2 Modic change.)]

Studies to date report relief of pain and improvement in function and quality of life after treatment, however, most are company sponsored, limited in size and are of generally poor or fair quality. Long-term non-industry-funded prospective trials should be pursued to confirm the results of currently published clinical studies.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2020, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
64400	Injection(s), anesthetic agent(s) and/or steroid; trigeminal nerve, each branch (ie, ophthalmic, maxillary, mandibular)
64405	Injection(s), anesthetic agent(s) and/or steroid; greater occipital nerve
64408	Injection(s), anesthetic agent(s) and/or steroid; vagus nerve

CPT[®] Codes	Description
64415	Injection(s), anesthetic agent(s) and/or steroid; brachial plexus
64417	Injection(s), anesthetic agent(s) and/or steroid; axillary nerve
64418	Injection(s), anesthetic agent(s) and/or steroid; suprascapular nerve
64420	Injection(s), anesthetic agent(s) and/or steroid; intercostal nerve, single level
64421	Injection(s), anesthetic agent(s) and/or steroid; intercostal nerve, each additional level
64425	Injection(s), anesthetic agent(s) and/or steroid; ilioinguinal, iliohypogastric nerves
64430	Injection(s), anesthetic agent(s) and/or steroid; pudendal nerve
64435	Injection(s), anesthetic agent(s) and/or steroid; paracervical (uterine) nerve
64445	Injection(s), anesthetic agent(s) and/or steroid; sciatic nerve
64447	Injection(s), anesthetic agent(s); femoral nerve
64450	Injection(s), anesthetic agent(s) and/or steroid; other peripheral nerve or branch
64454	Injection(s), anesthetic agent(s) and/or steroid; genicular nerve branches, including imaging guidance, when performed
64505	Injection, anesthetic agent; sphenopalatine ganglion
64510	Injection, anesthetic agent; stellate ganglion (cervical sympathetic)
64517	Injection, anesthetic agent; superior hypogastric plexus
64520	Injection, anesthetic agent; lumbar or thoracic (paravertebral sympathetic)
64530	Injection, anesthetic agent; celiac plexus, with or without radiologic monitoring
64620	Destruction by neurolytic agent, intercostal nerve
64624	Destruction by neurolytic agent, genicular nerve branches including imaging guidance, when performed
64640	Destruction by neurolytic agent; other peripheral nerve or branch
64680	Destruction by neurolytic agent, with or without radiologic monitoring; celiac plexus
64999	Unlisted procedure, nervous system

HCPCS Codes	Description
N/A	

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

+ Indicates a code requiring an additional character

ICD-10-CM Code	Description
C25.0-C25.9	Malignant neoplasm of pancreas
G44.85	Primary stabbing headache
G50.1	Atypical facial pain
G54.0-G54.9	Nerve root and plexus disorders
G56.40-G56.43	Causalgia of upper limb
G57.70-G57.73	Causalgia of lower limb
G89.22	Chronic post-thoracotomy pain
G89.4	Chronic pain syndrome
G90.50-G90.59	Complex regional pain syndrome I (CRPS I)
M54.81	Occipital neuralgia

ICD-10-CM Code	Description
K86.0	Alcohol-induced chronic pancreatitis
K86.1	Other chronic pancreatitis
R07.81-R07.89	Other chest pain
R10.10-R10.12	Pain localized to upper abdomen
S22.41X+ S22.49X+	Multiple fractures of rib

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Policy split from CP.MP.118 Injections for Pain Management. Sympathetic nerve block for CRPS: reworded diagnostic criteria for CRPS, retaining clinical meaning; added requirement of positive response to first or second block if requesting additional; added that blocks should be at least one week apart. Expanded criteria for sympathetic nerve block for pancreatic cancer to include celiac plexus neurolysis and gave it its own section. Changed indication for ischemic leg pain from “limited evidence to support” to “not medically necessary.” Updated background. References reviewed and updated. Coding updated.	08/18	08/18
Annual review. References reviewed and updated (added International Headache Society and Practice Guidelines for Chronic Pain Management). Specialty review completed. Removed CPT 64508 as code was inactive 1/1/2019. Added CPT 64620 for intercostal neurolysis. Specified that the following codes DO NOT support medical necessity: 64400, 64402, 64408, 64410, 64413, 64415, 64417, 64418, 64425, 64430, 64435, 64445, 64447, 64450, 64505.	08/19	08/19
Peripheral/Ganglion Nerve Blocks: Section A indication added for peripheral nerve blocks for malignant pain; section B.1. and 2. added indication for diagnosis or treatment of post-herniorrhaphy pain and therapeutic post-herniorrhaphy pain; section C added peripheral nerve blocks for prevention or treatment of headaches, including migraines, refractory migraines in pregnancy, and short-lasting unilateral neuralgiform headaches as not medically necessary. Corrected V. on Genicular Nerve Blocks and Neurotomy to state that they are experimental vs not medically necessary. Background and references updated accordingly. Combined all CPT codes into one table.	10/19	10/19
Added “neurolysis” as a not medically necessary procedure to section V. on genicular nerve block. Removed CPT codes 64402, 64410, and 64413- codes deleted 1/1/20 and replaced with unlisted code, 64999 as directed per CPT manual. Revised description for the following CPT codes effective 1/20: 64400-64450. Added 2020 CPT codes 64454 and 64624.	01/20	02/20
For occipital nerve block, added “trigger point at the emergence of the greater occipital nerve or in the distribution of C2” as an alternative to	07/20	08/20

Reviews, Revisions, and Approvals	Revision Date	Approval Date
tenderness at the affected nerve branch. Revised examples of less invasive pain medication in VI.B.c., “NSAIDs and opiates” to “NSAIDs and/or opiates.” References reviewed and updated.		
Added reference to CP.PHAR 232 for requests for Botox for migraine. Replaced “member” with “member/enrollee.”	12/20	
Added the following note to VI. Peripheral/ganglion nerve blocks: Peripheral/ganglion nerve blocks may be approved without prior authorization when used during another medically necessary procedure (i.e. as anesthesia during surgery).	05/21	
Annual review. Added refractory chronic pancreatitis as an indication for celiac plexus block to section III and updated background accordingly. Added ICD -10 codes K86.0 & K86.1 to support coverage criteria. Changed “Experimental/investigational” language in section V. and VI.E. to “insufficient evidence to support...”. Under section VI, moved “Note” for visibility. Added insufficient evidence to support peripheral nerve block for treatment of trigeminal neuralgia to VI.D, removed G50.0 from list of ICD 10 codes that support coverage criteria and updated background accordingly. References reviewed, reformatted and updated. Changed “review date” in the header to “date of last revision” and “date” in the revision log header to “revision date.” Reviewed by specialist.	08/21	08/21
Edited note in section VI to state: If administered as part of a surgery or other procedure, coding for peripheral/ganglion nerve blocks should follow proper coding practices and would not be subject to prior authorization or payment separately from the procedure.	09/21	
Revised policy title from “Nerve Blocks for Pain Management” to “Nerve Blocks and Neurolysis for Pain Management.” Added VII. Insufficient evidence to determine the safety and effectiveness of intraosseous radiofrequency nerve ablation of basivertebral nerve. Updated background and references accordingly.	12/21	12/21

References

1. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia*. 2013;33(9):629-808. doi:10.1177/0333102413485658
2. Drewes AM, Campbell CM, Ceyhan GO, et al. Pain in pancreatic ductal adenocarcinoma: A multidisciplinary, International guideline for optimized management. *Pancreatology*. 2018;18(4):446-457. doi: 10.1016/j.pan.2018.04.008
3. Arcidiacono PG, Calori G, Carrara S, McNicol ED, Testoni PA. Celiac plexus block for pancreatic cancer pain in adults. *Cochrane Database Syst Rev*.2011;2011(3):CD007519. Published 2011 Mar 16. doi: 10.1002/14651858.CD007519.pub2
4. Bashir MM, Shahzad MA, Yousaf MN, Khan BA, Khan FA. Comparison of postoperative pain relief by intercostal block between pre-rib harvest and post-rib harvest groups. *J College Physicians Surg Pak*. 2014;24(1):43-46.

CLINICAL POLICY

Nerve Blocks

5. Sarani B. Inpatient management of traumatic rib fractures. UpToDate. www.uptodate.com. Published October 25, 2019. Accessed July 29, 2021.
6. Hwang EG, Lee Y. Effectiveness of intercostal nerve block for management of pain in rib fracture patients. *Exerc Rehabil*. 2014;10(4):241-24. doi:10.12965/jer.140137
7. Morimoto Y. (2019) Regional anesthesia for thoracic surgery. *Anesthesia Pain and Intensive Care*. 352-356. 2010-2013
8. Rice DC, Cata JP, Mena GE, Rodriguez-Restrepo A, Correa AM, Mehran RJ. Posterior Intercostal Nerve Block with Liposomal Bupivacaine: An Alternative to Thoracic Epidural Analgesia. *Ann Thorac Surg*. 2015;99(6):1953-1960.
9. Health Technology Assessment: Genicular Nerve Block for the Management of Knee Pain. Hayes. www.hayesinc.com. Published June 24, 2020. (annual review July 1, 2021.) Accessed July 29, 2021.
10. McCormick ZL, Reddy R, Korn M, et al. A Prospective Randomized Trial of Prognostic Genicular Nerve Blocks to Determine the Predictive Value for the Outcome of Cooled Radiofrequency Ablation for Chronic Knee Pain Due to Osteoarthritis. *Pain Med*. 2018;19(8):1628-1638.
11. Kesikburun S, YaSar E, Uran A, Adiguzel E, Yimaz B. Ultrasound-guided genicular nerve pulsed radiofrequency treatment for painful knee osteoarthritis: A preliminary report. *Pain Physician* 2016;19(5):E751-E759.
12. Qudsi-Sinclair S, Borrás-Rubio E, Abellan-Guillén JF, Padilla Del Rey ML, Ruiz-Merino G. A Comparison of Genicular Nerve Treatment Using Either Radiofrequency or Analgesic Block with Corticosteroid for Pain after a Total Knee Arthroplasty: A Double-Blind, Randomized Clinical Study. *Pain Pract*. 2017;17(5):578-588. doi:10.1111/papr.12481
13. Ahmed A, Arora D. Ultrasound-guided radiofrequency ablation of genicular nerves of knee for relief of intractable pain from knee osteoarthritis: a case series. *Br J Pain*. 2018;12(3):145-154. doi: 10.1177/2049463717730433
14. Kim DH, Choi SS, Yoon SH, et al. Ultrasound-Guided Genicular Nerve Block for Knee Osteoarthritis: A Double-Blind, Randomized Controlled Trial of Local Anesthetic Alone or in Combination with Corticosteroid. *Pain Physician*. 2018;21(1):41-52.
15. Choi WJ, Hwang SJ, Song JG, et al. Radiofrequency treatment relieves chronic knee osteoarthritis pain: a double-blind randomized controlled trial. *Pain*. 2011;152(3):481-487. doi: 10.1016/j.pain.2010.09.029
16. Kucera TJ, Boezaart AP. Regional anesthesia does not consistently block ischemic pain: two further cases and a review of the literature. *Pain Med*. 2014;15(2):316-319. doi: 10.1111/pme.12235. Epub 2013 Sep 18.
17. McGreevy K, Hurley RW, Erdek MA, Aher MM, Li S, Cohen SP. The effectiveness of repeat celiac plexus neurolysis for pancreatic cancer: a pilot study. *Pain Pract*. 2013;13: 89–95. doi:10.1111/j.1533-2500.2012.00557.x
18. Gonzalez Sotelo V, Maculée F, Minguell J, Bergé R, Franco C, Sala-Blanch X... Ultrasound-guided genicular nerve block for pain control after total knee replacement: Preliminary case series and technical note. *Rev Esp Anesthesiol Reanim*.2017;64(10):568-576. doi:10.1016/j.redar.2017.04.001
19. Garza I. Occipital neuralgia. UpToDate. www.uptodate.com. Published February 22, 2021. Accessed July 29, 2021.
20. Rosenblatt MA, Lai Y. Scalp block and cervical plexus block techniques. UpToDate. www.uptodate.com. Published July 22, 2021. Accessed July 30, 2021.

21. Abdi S. Complex regional pain syndrome in adults: Pathogenesis, clinical manifestations and diagnosis. UpToDate. www.uptodate.com. Published July 13, 2020. Accessed August 2, 2021.
22. Abdi S. Complex regional pain syndrome in adults: Treatment, prognosis and prevention. UpToDate. www.uptodate.com. Published July 14, 2020. Accessed August 2, 2021.
23. Fernandez-del Castillo C, Jimenez RE, Murphy JE. Supportive care of the patient with locally advanced or metastatic exocrine pancreatic cancer. UpToDate. www.uptodate.com. Published April 02, 2021. Accessed August 2, 2021.
24. Freedman SD, Forsmark CE. Chronic pancreatitis: Management. UpToDate, Published June 22, 2021. Accessed Aug 2, 2021.
25. Health Technology Assessment: Local Injection Therapy for Cervicogenic Headache and Occipital Neuralgia. Hayes. www.hayesinc.com. Published September 28, 2017. (annual review January 29, 2021) Accessed July 29, 2021.
26. Soloman M, Mekhail MN, Mekhail N. Radiofrequency treatment in chronic pain. *Expert Rev Neurother*. 2010;10(3):469-474. doi:10.1586/ern.09.153
27. Portenoy RK, Copenhaver DJ. Cancer pain management: Interventional therapies. UpToDate. www.uptodate.com. Published May 19, 2020. Accessed August 2, 2021.
28. Lavu H, Lengel HB, Sell NM, et al. A prospective, randomized, double-blind, placebo controlled trial on the efficacy of ethanol celiac plexus neurolysis in patients with operable pancreatic and periampullary adenocarcinoma. *J Am Coll Surg*. 2015;220(4):497-508. doi: 10.1016/j.jamcollsurg.2014.12.013
29. Wyse JM, Carone M, Paquin SC, Usatii M, Sahai AV. Randomized, double-blind, controlled trial of early endoscopic ultrasound-guided celiac plexus neurolysis to prevent pain progression in patients with newly diagnosed, painful, inoperable pancreatic cancer. *J Clin Oncol*. 2011;29(26):3541-3546. doi:10.1200/JCO.2010.32.2750
30. American Society of Anesthesiologists Task Force on Chronic Pain Management; American Society of Regional Anesthesia and Pain Medicine. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology* 2010;112(4):810-833. doi: 10.1097/ALN.0b013e3181c43103
31. Lee MJ, Guinn D, Hickenbottom, S. Headache in pregnant and postpartum women. UpToDate. www.uptodate.com. Published Dec 2, 2020. Accessed August 2, 2021.
32. Matharu MS, Cohen AS. Short-lasting unilateral neuralgiform headache attacks: Treatment and prognosis. UpToDate. www.uptodate.com. Published June 21, 2021. Accessed August 2, 2021.
33. Tauben D, Stacey BR. Approach to the management of Chronic Non-cancer Pain in adults. UpToDate. www.uptodate.com. Published November 20, 2020. Accessed July 30, 2021.
34. Smith TJ, Saiki CB. Cancer Pain Management. *Mayo Clin Proc*. 2015;90(10):1428-1439. doi: 10.1016/j.mayocp.2015.08.009
35. Chambers WA. Nerve blocks in palliative care. *Br J Anaesth*. 2008; P 101(1):95-100. doi:10.1093/bja/aen105
36. Bonwich, MD, JB. Post-herniorrhaphy Groin Pain. UpToDate. www.uptodate.com. Published March 02, 2020. Accessed July 30, 2021.
37. Jeng, CL, Rosenblatt, MA. Overview of Peripheral Nerve Blocks. UpToDate. www.uptodate.com. Published October 20, 2020. Accessed July 30, 2021.

CLINICAL POLICY**Nerve Blocks**

38. Phillips, K, Schur PH. Management of isolated musculoskeletal chest pain. UpToDate. www.uptodate.com. Published May 20, 2020. Accessed August 2, 2021.
39. Levy MJ, Wiersema, MJ. Endoscopic ultrasound-guided celiac plexus interventions for pain related to pancreatic disease. UpToDate. www.uptodate.com. Published February 25, 2021.
40. Health Technology Assessment: Radiofrequency Nerve Ablation for the Management of Osteoarthritis of the Knee. Hayes. www.hayesinc.com. Published December 22, 2020. Accessed July 29, 2021.
41. Gardner TB, Adler DG, Forsmark CE, Sauer BG, Taylor JR, Whitcomb DC. ACG Clinical Guideline: Chronic Pancreatitis. *Am J Gastroenterol*. 2020;115(3):322-339. doi:10.14309/ajg.0000000000000535
42. Ho CC, Khan SA, Whealy MA. Trigeminal neuralgia. UpToDate. www.uptodate.com. Published August 3, 2020. Accessed August 4, 2021.
43. Michalik A, Conger A, Smuck M, Maus TP, McCormick ZL. Intraosseous Basivertebral Nerve Radiofrequency Ablation for the Treatment of Vertebral Body Endplate Low Back Pain: Current Evidence and Future Directions. *Pain Med*. 2021;22(Suppl 1):S24-S30. doi:10.1093/pm/pnab117
44. Evolving Evidence Review. Intracept Intraosseous Nerve Ablation System (Relievent Medsystems Inc.) for Treatment of Adults With Low Back Pain. Hayes. www.hayesinc.com. Published July 30, 2021. Accessed December 6, 2021.
45. Fischgrund JS, Rhyne A, Franke J, et al. Intraosseous basivertebral nerve ablation for the treatment of chronic low back pain: a prospective randomized double-blind sham-controlled multi-center study. *Eur Spine J*. 2018;27(5):1146-1156. doi:10.1007/s00586-018-5496-1
46. Fischgrund JS, Rhyne A, Macadaeg K, et al. Long-term outcomes following intraosseous basivertebral nerve ablation for the treatment of chronic low back pain: 5-year treatment arm results from a prospective randomized double-blind sham-controlled multi-center study. *Eur Spine J*. 2020;29(8):1925-1934. doi:10.1007/s00586-020-06448-x
47. Khalil JG, Smuck M, Koreckij T, et al. A prospective, randomized, multicenter study of intraosseous basivertebral nerve ablation for the treatment of chronic low back pain. *Spine J*. 2019;19(10):1620-1632. doi:10.1016/j.spinee.20
48. Fischgrund JS, Rhyne A, Franke J, et al. Intraosseous Basivertebral Nerve Ablation for the Treatment of Chronic Low Back Pain: 2-Year Results From a Prospective Randomized Double-Blind Sham-Controlled Multicenter Study. *Int J Spine Surg*. 2019;13(2):110-119. Published 2019 Apr 30. doi:10.14444/601519.05.598
49. 510(k) Premarket Notification. Intracept Intraosseous Nerve Ablation System (RF Probe), Intracept Intraosseous Nerve Ablation System (Access Instruments), Relievent RF Generator. Summary of Safety and Effectiveness. U.S. Food and Drug Administration Center for Devices and Radiological Health Web site. https://www.accessdata.fda.gov/cdrh_docs/pdf19/K190504.pdf. Published May 3, 2019. Accessed December 6, 2021.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical

policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of member/enrollees. This clinical policy is not intended to recommend treatment for member/enrollees. Member/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, member/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, member/enrollees and their representatives agree to be bound by such terms and conditions by providing services to member/enrollees and/or submitting claims for payment for such services.

Note: For Medicaid member/enrollees, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

CLINICAL POLICY

Nerve Blocks

Note: For Medicare member/enrollees, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.